

**Table SP41.1B Methamphetamine Use in the Past Year among Persons Aged 12 or Older, by State and Age Group: Percentages, Annual Averages
Based on 2004-2007**

State	Past Year Methamphetamine Use ¹				State	Past Year Methamphetamine Use ¹			
	Age Group					Age Group			
	Total	12-17	18-25	26 or Older		Total	12-17	18-25	26 or Older
Total United States	0.68	0.65	1.65	0.51	Missouri	0.80	0.95	2.50	0.49
Alabama	0.81	0.99	1.89	0.60	Montana	1.15	1.35	3.01	0.79
Alaska	0.94	0.52	3.09	0.60	Nebraska	1.01	0.50	2.53	0.78
Arizona	1.58	0.90	3.71	1.29	Nevada	1.63	1.33	3.97	1.31
Arkansas	1.84	1.42	4.23	1.48	New Hampshire	0.19	0.42	0.38	0.13
California	1.34	1.14	2.78	1.10	New Jersey	0.07	0.17	0.44	*
Colorado	1.43	0.84	2.15	1.39	New Mexico	1.67	0.82	3.80	1.38
Connecticut	0.10	0.11	0.54	0.03	New York	0.19	0.14	0.33	0.17
Delaware	0.29	0.42	0.88	0.17	North Carolina	0.32	0.33	0.58	0.27
District of Columbia	0.29	0.04	0.59	0.26	North Dakota	0.80	1.15	1.11	0.69
Florida	0.35	0.34	1.18	0.23	Ohio	0.29	0.49	0.98	0.14
Georgia	0.81	0.99	1.19	0.72	Oklahoma	1.14	0.68	3.03	0.85
Hawaii	1.01	0.67	1.54	0.96	Oregon	1.23	0.74	3.47	0.92
Idaho	1.38	1.59	3.74	0.89	Pennsylvania	0.26	0.36	0.48	0.21
Illinois	0.27	0.28	0.41	0.24	Rhode Island	0.24	0.51	0.32	0.19
Indiana	0.61	0.65	1.66	0.42	South Carolina	0.56	0.49	2.29	0.28
Iowa	0.82	1.11	1.90	0.58	South Dakota	1.09	1.27	2.75	0.74
Kansas	1.07	1.02	1.84	0.93	Tennessee	0.39	0.30	1.48	0.23
Kentucky	1.03	0.67	1.65	0.97	Texas	0.74	0.92	2.07	0.45
Louisiana	0.70	0.37	1.93	0.51	Utah	1.10	0.49	1.64	1.06
Maine	0.38	1.36	1.10	0.15	Vermont	0.11	0.17	0.68	*
Maryland	0.17	0.28	0.40	0.11	Virginia	0.37	0.55	0.96	0.25
Massachusetts	0.13	0.41	0.23	0.08	Washington	1.45	0.93	2.90	1.27
Michigan	0.35	0.43	1.04	0.23	West Virginia	0.58	0.38	1.94	0.39
Minnesota	0.65	0.59	3.06	0.22	Wisconsin	0.56	0.60	1.36	0.41
Mississippi	0.59	0.49	1.79	0.37	Wyoming	1.62	1.81	4.61	1.03

*Low precision; no estimate reported.

¹ Estimates of Past Year Methamphetamine Use include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2006 data year. For the 2004 and 2005 data, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with the 2006 and 2007 data. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.